

OptiPrep™ Application Sheet S41

Resolution of smooth endoplasmic reticulum (SER) and SER domains

- ◆ OptiPrep™ is a 60% (w/v) solution of iodixanol in water, density = 1.32 g/ml
- ◆ To access other Application Sheets referred to in the text return to the Subcellular Membranes Index; key Ctrl “F” and type the S-Number in the Find Box.

1. Background

It is widely recognized that the smooth endoplasmic reticulum contains specialized domains that are structurally and functionally associated with other membrane compartments. [Application Sheet S22](#) describes the fractionation of endoplasmic reticulum (ER), Golgi, plasma membrane and endosomes in continuous gradients of iodixanol, using relatively low *g*-forces for at least 12 h. The method is widely used and it highlights a paper by Woods et al [1], which describes this strategy as applied to 3T3 cells. Calreticulin shows a distinctive biphasic distribution in a 10–40% (w/v) iodixanol gradient but only the denser fraction also contains paxillin, which identifies this ER subfraction as perinuclear [1]. This functional and structural specialization of the ER is now widely recognized as more and more functional specializations have been discovered. Lynes and Simmen [2], amongst others, have reviewed some of these domain-specific functions; for example the peripheral ER that is closely associated with the plasma membrane, the mitochondria-associated membranes (MAM) and domains of the smooth ER that are associated with peroxisome biogenesis and the formation of lipid droplets.

- ◆ This Application Sheet summarizes some of the iodixanol gradient methods that have been used for the study of smooth ER subfractions and the resolution of lipid droplets.

2. MAM analysis

Lewin et al [3] prepared a standard rat liver homogenate (see [Application Sheet S05](#)) and loaded a post-nuclear supernatant (PNS) on to 20–40% iodixanol gradient. No other details about the gradient were given. The principal aim of their studies was to determine the localization of acyl-CoA synthetase 4. The gradient fractions were analyzed for wide range of markers including: acyl-CoA synthetase 1 (ER located); mitochondria were identified by acyl-CoA synthetase 5, glutamate dehydrogenase and glycerol-3-phosphate acyltransferase. MAM was identified by phosphatidylethanolamine methyltransferase. The gradient was able to resolve peroxisomes, mitochondria, MAM and endoplasmic reticulum. An approximate indication of the distribution of the major membrane compartments that were analyzed is given in Figure 1.




Figure 1. Approx. distribution of rat liver membranes in 20–40% iodixanol gradient – data adapted from ref 3.

In studies by Myhill et al [4], HeLa cells were homogenized in a routine HEPES- buffered 0.25 M sucrose solution containing 1 mM EDTA, using a ball-bearing homogenizer. A PNS was loaded on to a continuous 5–25% (w/v) iodixanol gradient (produced by diffusion from a 5%-interval step gradient) at approx. 120,000 *g* for 3 h. An approximate indication of the distribution of the major membrane compartments that were analyzed is given in Figure 2. A very noticeable difference between the two patterns is the relative banding positions of MAM to the ER and mitochondria. In the case of rat liver the MAM was well resolved from the mitochondria but overlapped the ER, while in the case of HeLa cells the MAM were well resolved from the ER but co-banded with the mitochondria. It is not clear if this is




Figure 2. Approx. distribution of HeLa cell membranes in 5–25% iodixanol gradient – data adapted from ref 4.

a distinction between the two sources or the difference in the density range of the iodixanol gradient, or both. The 20-40% (w/v) iodixanol gradient used by Lewin et al [3] covers the range 1.127-1.223 g/ml while that used by Myhill et al [4] was 1.054-1.151 g/ml. The latter gradient has also been used to study (a) the localization of the ER oxidoreductin (Ero1 α) to MAM in HEK cells, which was found to be dependent on the oxidizing conditions in the ER [5] and (b) that in HeLa cells Rab32 regulates the properties of MAM, notably Ca²⁺ and the enrichment of calnexin [6]. Discontinuous gradients of 10-30% (w/v) iodixanol, similar to those used by Myhill et al [4] and Gilady et al [5] also resolved Golgi, ER and MAM [7] and revealed that palmitoylation of the trans-membrane thioredoxin family protein (TMX) and calnexin influence their enrichment in MAM. Moreover the palmitoylation of calnexin influenced its functional properties [8].

- ◆ Reviews of some of the methodology for the study of MAM is provided in refs 9 and 10.
- ◆ Iodixanol gradients have also been used for the clear resolution of SER, principally from lysosomes, but also other organelles such as peroxisomes and mitochondria, subsequent to an initial sucrose gradient fractionation. The methodology, developed by Radhakrishnan et al for CHO-K1 cells [11,12], has been extended to HEK cells [13-15], Niemann-Pick type C cells [16], HepG2 cells [17], HeLa cells [18] and mouse liver [19]

3. Lipid droplets

Lipid droplets have been observed to be associated with the ER for over thirty years but it is only relatively recently that they have been shown to be involved in viral infection and a number of lipid-associated diseases. Presently, rather few papers have been published in which an OptiPrep™-based method has been used in their isolation. Because of the growing interest in these particles however a short summary of the methodology is included here. The most complete information comes from a paper by Heid et al [20]. Human hepatocellular carcinoma cells were homogenized by nitrogen cavitation and the PNS was adjusted to 30% (w/v) iodixanol. It was overlaid with layers of 20% and 10% iodixanol and centrifuged either at 190,000 g for 3 h or 220,000g for 2 h. The lipid droplets were concentrated close to the top of the gradient. The same three layer flotation gradient was adopted by Suzuki et al [21] for HeLa cells and by Akil et al [22] for Huh7 cells; the centrifugation conditions were however rather different: 166,000 g for 5 h and 200,000 g for 16 h respectively. A similar flotation approach was used by Buers et al [23] for studies on macrophages, the gradient however spanned a higher density range, it comprised 40%, 30% and 20% (w/v) iodixanol, but a much reduced g-force of 10,000 g for only 1h. In all cases the lipid droplets were recovered from the top of the gradient.

4. References

1. Woods, A.J., Roberts, M.S., Choudhary, J., Barry, S.T., Mazaki, Y., Sabe, H., Morley, S.J., Critchley, D.R. and Norman, J.C. (2002) *Paxillin associates with poly(A)-binding protein 1 at the dense endoplasmic reticulum and the leading edge of migrating cells* J. Biol. Chem., **277**, 6428-6437
2. Lynes, E.M. and Simmen, T. (2011) *Urban planning of the endoplasmic reticulum (ER): How diverse mechanisms segregate the many functions of the ER* Biochim. Biophys. Acta, **1813**, 1893–1905
3. Lewin T.M., Van Horn, C.G., Krisans, S.K. and Coleman, R.A. (2002) *Rat liver acyl-CoA synthetase 4 is a peripheral-membrane protein located in two distinct subcellular organelles, peroxisomes and mitochondrial-associated membranes* Arch. Biochem. Biophys., **404**, 263-270
4. Myhill, N., Lynes, E.M., Nanji, J.A., Blagoveshchenskaya, A.D., Fei, H., Simmen, K.C., Cooper, T.J., Thomas, G. and Simmen, T. (2008) *The subcellular distribution of calnexin is mediated by PACS-2* Mol. Biol. Cell, **19**, 2777-2788
5. Gilady, S.Y., Bui, M., Lynes, E.M., Benson, M.D., Watts, R., Vance, J.E. and Simmen, T. (2010) *Ero1 α requires oxidizing and normoxic conditions to localize to the mitochondria-associated membrane (MAM)* Cell Stress Chaperones, **15**, 619–629
6. Bui, M., Gilady, S.Y., Fitzsimmons, R.E.B., Benson, M.D., Lynes, E.M., Gesson, K., Alto, N.M., Strack, S., Scott, J.D. and Simmen, T. (2010) *Rab32 modulates apoptosis onset and mitochondria-associated membrane (MAM) properties* J. Biol. Chem., **285**, 31590–31602
7. Lynes, E.M., Bui, M., Yap, M.C., Benson, M.D., Schneider, B., Ellgaard, L., Berthiaume, L.G. and Simmen, T. (2012) *Palmitoylated TMX and calnexin target to the mitochondria-associated membrane* EMBO J., **31**, 457–470

8. Lynes, E.M., Raturi, A., Shenkman, M., Sandoval, C.O., Yap, M.C., Wu, J., Janowicz, A., Myhill, N., Benson, M.D., Campbell, R.E., Berthiaume, L.G., Lederkremer, G.Z. and Simmen, T. (2013) *Palmitoylation is the switch that assigns calnexin to quality control or ER Ca²⁺ signaling* J. Cell Sci., **126**, 3893–3903
9. Marchi, S., Paternani, S. and Pinton, P. (2014) *The endoplasmic reticulum–mitochondria connection: One touch, multiple functions* Biochim. Biophys. Acta, **1837**, 461–469
10. Giacomello, M. and Pellegrini, L. (2016) *The coming of age of the mitochondria–ER contact: a matter of thickness* Cell Death Differentiat., **23**, 1417–1427
11. Radhakrishnan, A., Goldstein, J.L., McDonald, J.G. and Brown, M.S. (2008) *Switch-like control of SREBP-2 transport triggered by small changes in ER cholesterol: a delicate balance* Cell Metab., **8**, 512–521
12. Sokolov, A. and Radhakrishnan, A. (2010) *Accessibility of cholesterol in endoplasmic reticulum membranes and activation of SREBP-2 switch abruptly at a common cholesterol threshold* J. Biol. Chem., **285**, 29480–29490
13. Harrison, K.D., Miao, R.Q., Fernandez-Hernández, C., Suárez, Y., Dávalos, A. and Sessa, W.C. (2009) *Nogo-B receptor stabilizes Niemann-Pick type C2 protein and regulates intracellular cholesterol trafficking* Cell Metab., **10**, 208–218
14. Alexia, C., Poalas, K., Carvalho, G., Zemirli, N., Dwyer, J., Dubois, S.M., Hatchi, E.M., Cordeiro, N., Smith, S.S., Castanier, C., Le Guelte, A., Wan, L., Kang, Y., Vazquez, A., Gavard, J., Arnoult, D. and Bidère, N. (2013) *The endoplasmic reticulum acts as a platform for ubiquitinated components of nuclear factor κB signaling* Sci. Signal., **6(291)**, ra79
15. Kawaguchi, K., Okamoto, T., Morita, M. and Imanaka, T. (2016) *Translocation of the ABC transporter ABCD4 from the endoplasmic reticulum to lysosomes requires the escort protein LMBD1* Sci. Rep., **6**: 30183
16. Abi-Mosleh, L., Infante, R.E., Radhakrishnan, A., Goldstein, J.L. and Brown, M.S. (2009) *Cyclodextrin overcomes deficient lysosome-to-endoplasmic reticulum transport of cholesterol in Niemann-Pick type C cells* Proc. Natl. Acad. Sci., **106**, 19316–19321
17. Wang, Y., Lam, W., Chen, S-R., Guan, F-L., Dutchman, G.E., Francis, S., Baker, D.C. and Cheng, Y-C. (2016) *Tylophorine analog DCB-3503 inhibited cyclin D1 translation through allosteric regulation of heat shock cognate protein 70* Sci. Rep., **6**: 32832
18. Ferencz, C-M., Guigas, G., Veres, A., Neumann, B., Stemmann, O. and Weiss, M. (2016) *Shaping the endoplasmic reticulum in vitro* Biochim. Biophys. Acta, **1858**, 2035–2040
19. Hager, L., Li, L., Pun, H., Liu, L., Hossain, M.A., Maguire, G.F., Naples, M., Baker, C., Magomedova, L., Tam, J., Adeli, K., Cummins, C.L., Connelly, P.W. and Ng, D.S. (2012) *Lecithin:cholesterol acyltransferase deficiency protects against cholesterol-induced hepatic endoplasmic reticulum stress in mice* J. Biol. Chem., **287**, 20755–20768
20. Heid, H., Rickelt, S., Zimbelmann, R., Winter, S., Schumacher, H. and Dorflinger, Y. (2013) *Lipid droplets, perilipins and cytokeratins – unraveled liaisons in epithelium-derived cells* PLoS One, **8**: e63061
21. Suzuki, M., Murakami, T., Cheng, J., Kano, H., Fukata, M. and Fujimoto, T. (2015) *ELMOD2 is anchored to lipid droplets by palmitoylation and regulates adipocyte triglyceride lipase recruitment* Mol. Biol. Cell, **26**, 2333–2342
22. Akil, A., Peng, J., Omrane, M., Gondeau, C., Desterke, C., Marin, M., Tronchère, H., Taveneau, C., Sar, S. et al (2016) *Septin 9 induces lipid droplets growth by a phosphatidylinositol-5-phosphate and microtubule-dependent mechanism hijacked by HCV* Nat. Comm., **7**: 12203
23. Buers, I., Robenek, H., Lorkowski, S., Nitschke, Y., Severs, N.J. and Hofnagel, O. (2009) *TIP47, a lipid cargo protein involved in macrophage triglyceride metabolism* Arterioscler. Thromb. Vasc. Biol., **29**, 767–773