

OptiPrep™ Application Sheet C17

Purification and processing of animal (non-human) and avian spermatozoa

- ◆ OptiPrep™ is a 60% (w/v) solution of iodixanol in water, density = 1.32 g/ml
- ◆ **For human spermatozoa see Application Sheet C18**
- ◆ To access other Application Sheets referred to in the text return to the Cell Index; key Ctrl “F” and type the C-Number in the Find Box
- ◆ Section 2 describes the optimal method for the enrichment of viable spermatozoa from bovine semen using OptiPrep™.
- ◆ Sections 3-5 summarize some of the other published methods for sperm from rodents (3), equine sperm (4), and *Xenopus laevis* sperm and spermatids (5).
- ◆ Section 6 describes the use of OptiPrep™ for cryopreservation of sperm from a number of species.

1. Introduction

In ejaculates, viable spermatozoa of normal morphology are sometimes a very low percentage of the total cell population. This Application Sheet presents a detailed protocol for the recovery of a highly viable fraction of bovine spermatozoa for use in fertilization. The recommended strategy involves, in the first instance, adjustment of the density of a semen sample to approx. 1.17 g/ml. Ideally, two lower density solutions are then layered on top, so that the viable semen of normal morphology band at the interface between these two layers. These cells are thus completely separated from both the non-viable cells (and any soluble material released from partially broken cells), which all remain in the load zone, and any morphologically abnormal cells that band at the top of the least dense layer. This result is depicted in Figure 1.

Iodixanol is the gradient solute of choice: firstly, all of the solutions are easily prepared by dilution of OptiPrep™ with any buffered saline or a special diluent formulated for the maintenance of sperm viability, while Nycodenz® solutions must be prepared from Nycodenz® powder. Secondly, to raise the density of the ejaculate to approx 1.170 g/ml it is necessary to mix it with a high-density medium (usually >1.26 g/ml). Nycodenz® solutions are hyperosmotic above $\rho = 1.16$ g/ml, thus the seminal fluid would also become hyperosmotic. Consequently with Nycodenz® the semen has to be loaded at a lower density in the middle, or top of the gradient. This is not the case with iodixanol; OptiPrep™ or a dense solution prepared from OptiPrep™ and the chosen diluent, can be added to a raw ejaculate without increasing its osmolality (see Notes 1 and 2).

2. Purification of bovine spermatozoa

2a. Solutions required

- A. OptiPrep™ (shake gently before use)
- B. Diluent: Hanks Buffered Salt Solution (HBSS) or other suitable ambient temperature diluent such as Ruthin Diluent (RD); see also Note 2.

2b. Protocol

1. Assess a freshly taken ejaculate for viability and then mix with an equal volume of Solution A to raise its density to approx. 1.170 g/ml.
2. Prepare the two gradient solutions: 1.119 g/ml (4 vol. OptiPrep™ + 7 vol. HBSS) and 1.154 g/ml (9 vol. OptiPrep™ + 10 vol. HBSS). Using RD mix respectively 4 vol. + 8 vol. and 9 vol. + 11 vol. (see Notes 2 and 3).
3. In a suitable tube (50 ml) layer 10 ml of each of the two gradient solutions and underlayer these with 10-15 ml of the sample-OptiPrep™ mixture (density approximately 1.17 g/ml), to form a three-step gradient (see Figure 1 and Notes 4 and 5).

4. Centrifuge the gradient in a swinging-bucket rotor at 1500 g_{av} for 20 min at approx. 20°C.
5. After centrifugation, deformed sperm, cytoplasmic droplets, detached heads and tails band at the top of the gradient (A). Motile cells of normal morphology band at the 1.119/1.154 g/ml interface (B) while in the loading area a pellet (D) and some particulate material in suspension (C) contain immotile sperm (see Figure 1). The sperm cells from interface B can then be checked for viability and fertility and stored (see Notes 6-9).




Figure 1: Purification of normal motile sperm cells in a discontinuous iodixanol gradient

2c. Notes

1. Iodixanol is essentially a dimer of Nycodenz®, it therefore has approximately twice the molecular mass and solutions have half the osmolality of Nycodenz®
2. The diluent solution may be any solution thought appropriate by individual workers. The final density of the diluted fractions will depend upon the density of the diluent. Common physiological salt solutions, such as phosphate-buffered saline, Hanks Buffered Salt Solution (HBSS) or a more complex medium such as RPMI, have densities close to 1.005 - 1.006 g/ml. It may however be preferable to use a medium designed to preserve the viability and motility of sperm cells at room temperature. These so-called ambient temperature diluents (for example Ruthin Diluent) frequently contain polyhydric alcohols such as sorbitol and consequently have a slightly higher density ($\rho = 1.018$ g/ml). Details of the Ruthin Diluent can be obtained from Dr Stuart Revell, Genus Freezing Unit, Llanrhydd, Ruthin, Denbighshire, LL15 2UP, UK.
3. The volume of OptiPrep™ and medium required to prepare the density solutions will vary with the density of the medium. For more information about preparing density gradient solutions for mammalian cells see [Application Sheet C01](#).
4. For more information on preparation of discontinuous gradients see [Application Sheet C02](#).
5. More recently Garrett et al [1] simplified the gradient: the semen (diluted first with Eqcellsire™) was mixed with an equal volume of OptiPrep™; 8 ml of this suspension was then overlaid with 1 ml of a 1.15 g/ml solution and centrifuged at 1000 g for 15 min. The viable sperm banded at the top of the gradient.
6. The quality of the semen has been assessed by using membrane integrity as an indicator of general cell function and viability. The Osmotic Resistance Test (ORT) described by Revell and Mrode [2] and the fluorescent analysis method described by Harrison and Vickers [3] have been used to check membrane integrity. The motile band from the 1.119/1.154 g/ml interface shows over 95% viability by these tests, while the pelleted material and particulate material remaining in the loading layer are found to be 99% non-viable cells by these methods.
7. Routinely, the viable, motile spermatozoa are diluted in skimmed milk, glucose and glycerol, to provide 1.5×10^7 per A.I. straw and deep-frozen. When the straws are subsequently thawed and subjected to ORT, 74% of the sperm are still viable as judged by membrane integrity and activity.
8. Ejaculates from other species have shown similar but not identical banding characteristics: small changes to the precise densities of the layers may be required. The solutions and protocol in this Application Sheet will serve as a useful starting point from which adjustments to the final densities of the gradient layers can be made, to optimize the fractionation of material from other species.
9. This methodology has also been reported in ref 4.

3. Rodent sperm

Separation of viable and non-viable sperm cells was achieved on double layer iodixanol gradient of 15% and 24% (w/v) iodixanol; after centrifugation at 400 g for 20 min the viable sperm banded at the interface of the two iodixanol solutions [5-8].

4. Equine sperm

Stuhtmann et al [9] and Heutelbeck et al [10,11] used similar two-layer gradients for stallion sperm of 1.090 and 1.165 g/ml or 1.090 and 1.170 g/ml and commented that these gradients promoted better retention of morphology and progressive motility and that the gradient purification was more effective if carried out just after collection rather than immediately before cryopreservation.

5. *Xenopus laevis* sperm and spermatids

A discontinuous iodixanol gradient of 12%, 20% and 30% (w/v) iodixanol was used to separate the lighter spermatids from the denser mature semen [12].

6. Concentration of semen on to a dense cushion, prior to freezing

Compared with simple pelleting, sedimentation on to a dense cushion of iodixanol prior to cooling and freezing considerably improves the recovery and motility of viable sperm. The cushion is usually OptiPrep™ itself and the centrifugation conditions vary from 800 g for 10 min to 1000 g for 20 min. To harvest the band of spermatozoa, a narrow metal cannula is used to remove the liquids above and below the cells. The spermatozoa band is often described as a “pellet”; a “sharp band” is the more accurate description. The method has been widely used for equine sperm [13-33]; this has also been observed for gazelle semen [34], elephant semen [35-38], boar semen [39,40], donkey semen [41], buffalo semen [42], porcine semen [43], ram semen [44] and rodent semen [45,46].

Some more recent references reporting the use of OptiPrep™ are listed in Section 7: refs. # 47-53

Important technical note: It is not known if any of the published methods using the iodinated density gradient medium Nycodenz® can be translated directly to the use of OptiPrep™. Although solutions of the same % (w/v) will have the same density, the osmolality of iodixanol solutions will be approx. half those of Nycodenz®. Unlike those of Nycodenz®, iodixanol solutions can be made approx. isoosmotic with plasma at all densities.

7. References

- Garrett, L.J.A., Revell, S.G. and Leese, H.J. (2008) *Adenosine triphosphate production by bovine spermatozoa and its relationship to semen fertilizing ability* J. Androl., **29**, 449-458
- Revell, S.G. and Mrode, R.A. (1994) *An osmotic resistance test for bovine semen* Animal Reprod. Sci., **36**, 77-86
- Harrison, R.A.P. and Vickers, S.E. (1990) *Use of fluorescent probes to assess membrane integrity in mammalian spermatozoa* J. Reprod. Fert., **88**, 343-352
- Beer-Ljubić, B., Aladrović, J., Marenjak, T.S., Majić-Balić, I., Laškaj, R. and Milinković-Tur, S. (2012) *Biochemical properties of bull spermatozoa separated in iodixanol density solution* Res. Vet. Sci., **92**, 292-294
- Stein, K.K., Go, J.C., Primakoff, P. and Myles, D.G. (2005) *Defects in secretory pathway trafficking during sperm development in Adam2 knockout mice* Biol. Reprod., **73**, 1032-1038
- Stein, K.K., Go, J.C., Lane, W.S., Primakoff, P. and Myles, D.G. (2006) *Proteomic analysis of sperm regions that mediate sperm-egg interactions* Proteomics, **6**, 3533-3543
- Nishimura, H., Myles, D.G. and Primakoff, P. (2007) *Identification of an ADAM2-ADAM3 complex on the surface of mouse testicular germ cells and cauda epididymal sperm* J. Biol. Chem., **282**, 17900-17907
- Nishimura, H., Gupta, S., Myles, D.G. and Primakoff, P. (2011) *Characterization of mouse sperm TMEM190, a small transmembrane protein with the trefoil domain: evidence for co-localization with IZUMO1 and complex formation with other sperm proteins* Reproduction, **141**, 437-451
- Stuhtmann, G., Oldenhof, H., Peters, P., Klewitz, J., Martinsson, G. and Sieme, H. (2012) *Iodixanol density gradient centrifugation for selecting stallion sperm for cold storage and cryopreservation* Anim. Reprod. Sci., **133**, 184-190
- Heutelbeck, A., Oldenhof, H., Henke, S., Martinsson, G. and Sieme, H. (2012) *Delayed cryopreservation of stallion sperm: effect of iodixanol density gradient centrifugation* J. Equine Vet. Sci., **32**, 488-489
- Heutelbeck, A., Oldenhof, H., Rohn, K., Martinsson, G., Morrell, J.M. and Sieme, H. (2015) *Use of density centrifugation for delayed cryopreservation of stallion sperm: perform sperm selection directly after collection or after storage?* Reprod. Dom. Anim., **50**, 76-83

12. Teperek, M., Simeone, A., Gaggioli, V., Miyamoto, K., Allen, G.E., Erkek, S., Kwon, T., Marcotte, E.M., Zegerman, P. et al (2016) *Sperm is epigenetically programmed to regulate gene transcription in embryos* Genome Res., **26**, 1034-1046
13. Revell, S.G., Pettit, M.T. and Ford, T.C. (1997) *Use of centrifugation over iodixanol to reduce damage when processing stallion sperm for freezing*. Proc. Joint Meeting, Society for the Study of Fertility, Abstr. Series No. 92, 38 (abstr.)
14. Sieme, H., Knop, K. and Rath, D. (2006) *Effects of cushioned centrifugation on sperm quality in stallion semen stored cooled at 5°C for 24h, and stored cooled for 2h or 24h and then frozen* Animal Reprod. Sci., **94**, 99-103
15. Loomis, P.R. (2006) *Advanced methods for handling and preparation of stallion semen* Vet. Clin. Equine **22**, 663-676
16. Saragusty, J., Gacitua, H., Pettit, M.T. and Arav, A. (2007) *Directional freezing of equine semen in large volumes* Reprod. Dom. Anim., **42**, 610-615
17. Waite, J.A., Love, C.C., Brinsko, S.P., Teague, S.R., Salazar Jr. J.L., Mancill, S.S. and Varner, D.D. (2008) *Factors impacting equine sperm recovery rate and quality following cushioned centrifugation* Theriogenology, **70**, 704-714
18. Varner, D.D., Love, C.C., Brinsko, S.P., Blanchard, T.L., Hartman, D.L., Bliss, S.B., Carroll, B.S. and Eslick, M.C. DVMD (2008) *Semen processing for the subfertile stallion* J. Equine Vet. Sci., **28**, 677-685
19. Webb, G.W. Dean, M.M. (2009) *Effect of centrifugation technique on post-storage characteristics of stallion spermatozoa* J. Equine Vet. Sci., **29**, 675-680
20. Mari, G., Morganti, C.M., Rizzato, G., Mislei, B., Iacono, E. and Merlo, B. (2010) *Comparison of density gradient and simple centrifugation of equine spermatozoa: effect on fertility of an oligospermic-subfertile stallion* Animal Reprod. Sci., **121S**, S153–S154
21. Salazar Jr, J.L., Teague, S.R., Love, C.C., Brinsko, S.P., Blanchard, T.L. and Varner, D.D. (2011) *Effect of cryopreservation protocol on postthaw characteristics of stallion sperm* Theriogenology, **76**, 409–418
22. Hoogewijs, M., Morrell, J., Van Soom, A., Govaere, J., Johannisson, A., Piepers, S., De Schauwer, C., De Kruif, A. and S. De Vlieghe (2011) *Sperm selection using single layer centrifugation prior to cryopreservation can increase thawed sperm quality in stallions* Equine Vet. J., **43** (Suppl. 40), 35-41
23. Blanchard, T.L., Brinsko, S.P., Love, C.C., Vest, D.D., Berezowski, C.B., Wendt, K.M., Stich, K. and Varner, D.D. (2012) *Case study of processing and insemination techniques: attempts to improve fertility of an aged stallion with dilute semen of poor quality* J. Equine Vet. Sci., **32**, 5-11
24. Edmond, A.J., Brinsko, S.P., Love, C.C., Blanchard, T.L., Teague, S.R. and Varner, D.D. (2012) *Effect of centrifugal fractionation protocols on quality and recovery rate of equine sperm* Theriogenology, **77**, 959–966
25. Bliss, S.B., Voge, J.L., Hayden, S.S., Teague, S.R., Brinsko, S.P., Love, C.C., Blanchard, T.L. and Varner, D.D. (2012) *The impact of cushioned centrifugation protocols on semen quality of stallions* Theriogenology **77**, 1232–1239
26. Love, C.C., Blanchard, T.L., Varner, D.D., Brinsko, S.P., Voge, J., Bliss, S., Sudderth, K., Teague, S. and LaCaze, K. (2012) *Effect of daily semen centrifugation and resuspension on the longevity of equine sperm quality following cooled storage* Theriogenology, **77**, 1911–1917
27. Len, J.A., Beehan, D.P., Lyle, S.K. and Eilts, B.E. (2013) *Cushioned versus noncushioned centrifugation: Sperm recovery rate and integrity* Theriogenology, **80**, 648–653
28. Ponthier, J., Franck, T., Parrilla-Hernandez, S., Niesten, A., de la Rebiere, G., Sertejn, D. and Deleuze, S. (2014) *Concentration, activity and biochemical characterization of myeloperoxidase in fresh and post-thaw equine semen and their implication on freezability* Reprod. Dom. Anim., **49**, 285–291
29. Bradecamp, E.A. (2014) *Centrifugation of semen: cushion technique* In Equine Reproductive Procedures, (ed. Dascanio, J. and McCue, P.) JohnWiley & Sons, Inc., pp 429-432
30. Heutelbeck, A., Oldenhof, H., Rohn, K., Martinsson, G., Morrell, J.M. and Sieme, H. (2015) *Use of density centrifugation for delayed cryopreservation of stallion sperm: perform sperm selection directly after collection or after storage?* Reprod. Dom. Anim., **50**, 76–83
31. Stawicki, R.J., McDonnell, S.M., Giguère, S. and Turner, R.M. (2016) *Pregnancy outcomes using stallion epididymal sperm stored at 5 °C for 24 or 48 hours before harvest* Theriogenology, **85**, 698–702
32. Blommaert, D., Franck, T., Donnay, I., Lejeune, J-P., Dettleux, J. and Sertejn, D. (2016) *Substitution of egg yolk by a cyclodextrin-cholesterol complex allows a reduction of the glycerol concentration into the freezing medium of equine sperm* Cryobiology, **72**, 27-32
33. Voge, J., Varner, D.D., Blanchard, T.L., Meschini, M., Turner, C., Teague, S.R., Brinsko, S.P. and Love, C.C. (2016) *The effects of urine concentration, and cushion centrifugation to remove urine, on the quality of cool-stored stallion sperm* Theriogenology **86**, 1294–1298

34. Saragusty, J., Gacitua, H., King, R. and Arav, A. (2006) *Post-mortem semen cryopreservation and characterization in two different endangered gazelle species (Gazella gazella and Gazella dorcas) and one subspecies (Gazella gazelle acaiae)* Theriogenology, **66**, 775-784
35. Hermesa, R., Behr, B., Hildebrandt, T.B., Blottner, S., Sieg, B., Frenzel, A., Knieriem, A., Saragusty, J. and Rath, D. (2009) *Sperm sex-sorting in the Asian elephant (Elephas maximus)* Animal Reprod. Sci., **112**, 390–396
36. Saragusty, J., Hildebrandt, T.B., Behr, B., Knieriem, A., Kruse, J. and Hermes, R. (2009) *Successful cryopreservation of Asian elephant (Elephas maximus) spermatozoa* Anim. Reprod. Sci., **115**, 255–266
37. Hildebrandt, T.B., Hermes, R., Saragusty, J., Potier, R., Schwammer, H.M., Balfanz, F., Vielgrader, H., Baker, B., Bartels, P. and Göritz, F. (2012) *Enriching the captive elephant population genetic pool through artificial insemination with frozen-thawed semen collected in the wild* Theriogenology, **78**, 1398–1404
38. Hermes, R., Saragusty, J., Goritz, F., Bartels, P., Potier, R., Baker, B., Streich, W.J. and Hildebrandt, T.B. (2013) *Freezing African elephant semen as a new population management tool* PLoS One **8**: e57616
39. Matás, C., Decuadro, G., Martínez-Miró, S. and Gadea, J. (2007) *Evaluation of a cushioned method for centrifugation and processing for freezing boar semen* Theriogenology **67**, 1087-1091
40. Zhang, W., Yi, K., Chen, C., Hou, X. and Zhou, X. (2012) *Application of antioxidants and centrifugation for cryopreservation of boar spermatozoa* Anim. Reprod. Sci., **132**, 123– 128
41. Saragusty, J., Lemma, A., Hildebrandt, T.B. and Goëritz, F. (2017) *Follicular size predicts success in artificial insemination with frozen-thawed sperm in donkeys* PLoS One, **12**: e0175637
42. Swami, D.S., Kumar, P., Malik, R.K., Saini, M., Kumar, D. and Jan, M.H. (2017) *The cryoprotective effect of iodixanol in buffalo semen cryopreservation* Anim. Reprod. Sci., **179**, 20–26
43. Romar, R., Funahashi, H. and Coy, P. (2016) *In vitro fertilization in pigs: New molecules and protocols to consider in the forthcoming years* Theriogenology, **85**, 125–134
44. Cirit, U., Bagis, H., Demir, K., Agca, C., Pabuccuoglu, S., Varisli, O., Clifford-Rathert, C., Agcac, Y. (2013) *Comparison of cryoprotective effects of iodixanol, trehalose and cysteamine on ram semen* Anim. Reprod. Sci., **139**, 38– 44
45. Kim, S., Agca, C. and Agca, Y. (2013) *Effects of iodixanol during rat epididymal sperm cryopreservation* Cryobiology, **67**, 398–442
46. Kim, S., Hooper, S., Agca, C. and Agca, Y. (2016) *Post-thaw ATP supplementation enhances cryoprotective effect of iodixanol in rat spermatozoa* Reprod. Biol. Endocrinol., **14**: 5
47. Chuawongboon, P., Sirisathien, S., Jatuporn Pongpeng, J., Sakhong, D., Nagai, T. and Vongpralub, T. (2017) *Effects of supplementation of iodixanol to semen extender on quality and fertilization ability of frozen-thawed Thai native bull sperm* Anim. Sci. J., **88**, 1311–1320
48. Ali, A., Ahmad, E., Ijaz, N., Ahmad, W. and Ulhassan, F. (2018) *Quality of cryopreserved buffalo spermatozoa improved from poor quality ejaculates* Cryobiology, **80**, 167
49. Agca, C., Timonin, M., Kim, S., Epperson, K. and Agca, Y. (2018) *Cryosurvival of Mus musculus and peromyscus spermatozoa in the presence of iodixanol* Cryobiology, **80**, 185
50. Brom-de-Luna, J.G., Siqueira Canesin, H., Wright, G. and Hinrichs, K. (2018) *Culture of somatic cells isolated from frozen-thawed equine semen using fluorescence-assisted cell sorting* Anim. Reprod. Sci., **190**, 10–17
51. Marqui, F.N., Martins Jr., A., da Cruz, T.E., Berton, T.I.U., de Paula Freitas-Dell’Aqua, C., Dell’Aqua Júnior, J.A., Oba, E. (2018) *Addition of iodixanol in bull freezing extender improves the sperm membranes integrity* Anim. Reprod. Sci. (Abstr), **194**, e1–e27
52. Hermes, R., Hildebrandt, T.B. and Göritz, F. (2018) *Cryopreservation in rhinoceros - setting a new benchmark for sperm cryosurvival* PLoS One, **13**: e0200154
53. Palazzese L., Gosalvez, J., Anzalone, D.A., Loi, P. and Saragusty, J. (2018) *DNA fragmentation in epididymal freeze-dried ram spermatozoa impairs embryo development* J. Reprod. Devel., **64**, 393-400